

## APPROXIMATION OF THE MINIMAL GERŠGORIN SET OF A SQUARE COMPLEX MATRIX\*

RICHARD S. VARGA<sup>†</sup>, LJILJANA CVETKOVIĆ<sup>‡</sup>, AND VLADIMIR KOSTIĆ<sup>‡</sup>

**Abstract.** In this paper, we address the problem of finding a numerical approximation to the minimal Geršgorin set,  $\Gamma^{\mathcal{R}}(A)$ , of an irreducible matrix  $A$  in  $\mathbb{C}^{n,n}$ . In particular, boundary points of  $\Gamma^{\mathcal{R}}(A)$  are related to a well-known result of Olga Taussky.

**Key words.** eigenvalue localization, Geršgorin theorem, minimal Geršgorin set.

**AMS subject classifications.** 15A18, 65F15

**1. Introduction.** Given an irreducible matrix  $A = [a_{i,j}]$  in  $\mathbb{C}^{n,n}$ , its  $i$ -th Geršgorin disk is defined, with  $N := \{1, 2, \dots, n\}$ , by

$$(1.1) \quad \Gamma_i(A) := \{z \in \mathbb{C} : |z - a_{i,i}| \leq r_i(A) := \sum_{j \in N \setminus \{i\}} |a_{i,j}|\} \quad (i \in N),$$

and the union of all these disks, denoted by

$$(1.2) \quad \Gamma(A) := \bigcup_{i \in N} \Gamma_i(A),$$

is called the *Geršgorin set* for  $A$ . A well-known result of Geršgorin [2] gives us that  $\Gamma(A)$  contains the spectrum,  $\sigma(A)$ , of  $A$ , i.e.,

$$(1.3) \quad \sigma(A) := \{\lambda \in \mathbb{C} : \det(\lambda I - A) = 0\} \subseteq \Gamma(A).$$

Continuing, for any  $\mathbf{x} = [x_1, x_2, \dots, x_n]^T > \mathbf{0}$  in  $\mathbb{R}^n$ , i.e.,  $x_i > 0$  for all  $i \in N$ , let  $X := \text{diag}[x_1, x_2, \dots, x_n]$  denote the associated nonsingular diagonal matrix. Then,  $X^{-1}AX$  has the same eigenvalues as  $A$ . Thus, with the Geršgorin disks for  $X^{-1}AX$  now given by

$$(1.4) \quad \Gamma_i^{\mathbf{x}}(A) := \{z \in \mathbb{C} : |z - a_{i,i}| \leq r_i^{\mathbf{x}}(A) := \sum_{j \in N \setminus \{i\}} \frac{|a_{i,j}|x_j}{x_i}\} \quad (i \in N),$$

and with the associated Geršgorin set,

$$(1.5) \quad \Gamma^{\mathbf{x}}(A) := \bigcup_{i \in N} \Gamma_i^{\mathbf{x}}(A),$$

then

$$(1.6) \quad \sigma(A) \subseteq \Gamma^{\mathbf{x}}(A), \quad \text{for any } \mathbf{x} > \mathbf{0} \text{ in } \mathbb{R}^n.$$

The inclusion of (1.6) is also a well-known result of Geršgorin [2]. Clearly, the following intersection,

$$(1.7) \quad \Gamma^{\mathcal{R}}(A) := \bigcap_{\mathbf{x} > \mathbf{0} \text{ in } \mathbb{R}^n} \Gamma^{\mathbf{x}}(A),$$

\*Received November 21, 2007. Accepted for publication October 27, 2008. Published online on December 17, 2008. Recommended by R. Nabben.

<sup>†</sup>Department of Mathematical Sciences, Kent State University, Kent, Ohio (varga@math.kent.edu).

<sup>‡</sup>Department of Mathematics and Informatics, Faculty of Science, University of Novi Sad, Serbia (lila,vkostic@im.ns.ac.yu).

called the *minimal Geršgorin set* in [4, 6], is always a subset of  $\Gamma^{r^x}(A)$ , for any  $\mathbf{x} > \mathbf{0}$  in  $\mathbb{R}^n$ , thereby giving the sharpest inclusion set for  $\sigma(A)$ , with respect to *all* positive diagonal similarity transforms  $X^{-1}AX$  of  $A$ .

This sharpness can also be expressed in the following way; cf. [6, Theorem 4.5]. With

$$(1.8) \quad \hat{\Omega}(A) := \{B = [b_{i,j}] \in \mathbb{C}^{n,n} : b_{i,i} = a_{i,i} \text{ and } |b_{i,j}| \leq |a_{i,j}| \text{ for } i \neq j (i, j \in N)\},$$

then

$$(1.9) \quad \sigma(\hat{\Omega}(A)) := \bigcup_{B \in \hat{\Omega}(A)} \sigma(B) = \Gamma^{\mathcal{R}}(A),$$

i.e., each point of  $\Gamma^{\mathcal{R}}(A)$  is an eigenvalue of *some* matrix  $B$  in  $\hat{\Omega}(A)$ .

Unlike the Geršgorin set  $\Gamma(A)$  of (1.2) or  $\Gamma^{r^x}(A)$  of (1.5), the minimal Geršgorin set  $\Gamma^{\mathcal{R}}(A)$  of (1.7) is not in general easy to determine numerically. The aim of this paper is to find a *reasonable approximation* of  $\Gamma^{\mathcal{R}}(A)$ , with a finite number of calculations, which contains  $\Gamma^{\mathcal{R}}(A)$ , and for which a limited number of boundary points of this approximation are actual boundary points of  $\Gamma^{\mathcal{R}}(A)$ . The determination of these latter boundary points are then related to a famous sharpening, by Olga Taussky [3], of the Geršgorin set of (1.2).

**2. Background.** Given an irreducible matrix  $A = [a_{i,j}]$  in  $\mathbb{C}^{n,n}$ , its associated irreducible matrix  $Q(z) = [q_{i,j}(z)]$ , in  $\mathbb{R}^{n,n}$ , is defined by

$$(2.1) \quad q_{i,i}(z) := -|z - a_{i,i}|, \text{ and } q_{i,j}(z) := |a_{i,j}|, \text{ for } i \neq j (i, j \in N).$$

If

$$(2.2) \quad \mu(z) := \max_{i \in N} |z - a_{i,i}|,$$

then the matrix  $B(z) := [b_{i,j}(z)] \in \mathbb{R}^{n,n}$ , defined by

$$(2.3) \quad b_{i,i}(z) := \mu(z) - |z - a_{i,i}|, \text{ and } b_{i,j}(z) := |a_{i,j}|, i \neq j (i, j \in N),$$

satisfies

$$(2.4) \quad B(z) = Q(z) + \mu(z)I_n,$$

where  $B(z)$  is a nonnegative irreducible matrix in  $\mathbb{R}^{n,n}$ . Then, from the Perron-Frobenius theory of nonnegative matrices, the matrix  $B(z)$  possesses a positive real eigenvalue,  $\rho(B(z))$ , called the *Perron root* of  $B(z)$ , which is characterized as follows. For any  $\mathbf{x} > \mathbf{0}$  in  $\mathbb{R}^{n,n}$ , either

$$(2.5) \quad \min_{i \in N} \{(B(z)\mathbf{x})_i / x_i\} < \rho(B(z)) < \max_{i \in N} \{(B(z)\mathbf{x})_i / x_i\},$$

or

$$(2.6) \quad B(z)\mathbf{x} = \rho(B(z))\mathbf{x}.$$

Thus, if we set

$$(2.7) \quad \nu(z) := \rho(B(z)) - \mu(z) \text{ (all } z \in \mathbb{C}),$$

then  $\nu(z)$  is a real-valued function, defined for all  $z \in \mathbb{C}$ . Moreover, from (2.5) and (2.6), for any  $\mathbf{x} > \mathbf{0}$  in  $\mathbb{R}^n$  and any  $z \in \mathbb{C}$ , either

$$(2.8) \quad \min_{i \in N} \{(Q(z)\mathbf{x})_i / x_i\} < \nu(z) < \max_{i \in N} \{(Q(z)\mathbf{x})_i / x_i\},$$

or

$$(2.9) \quad Q(z)\mathbf{x} = \nu(z)\mathbf{x},$$

the last equation giving us that  $\nu(z)$  is an eigenvalue of  $Q(z)$ .

The following connection of the function  $\nu(z)$  of (2.7) to the minimal Geršgorin set,  $\Gamma^{\mathcal{R}}(A)$ , comes from [4] and [6]:

$$(2.10) \quad z \in \Gamma^{\mathcal{R}}(A) \text{ if and only if } \nu(z) \geq 0,$$

and

$$(2.11) \quad \text{if } z \in \partial\Gamma^{\mathcal{R}}(A), \text{ then } \nu(z) = 0.$$

It is also known (cf. [6], Theorem 4.6), from the assumption that  $A$  is irreducible, that

$$(2.12) \quad \nu(a_{i,i}) > 0, \text{ for all } i \in N.$$

Further, given any real number  $\theta$  with  $0 \leq \theta < 2\pi$ , it is known (cf. [6], Theorem 4.6) that there is a largest number  $\hat{\rho}_i(\theta) > 0$  such that

$$(2.13) \quad \nu(a_{i,i} + \hat{\rho}_i(\theta)e^{i\theta}) = 0, \text{ and } \nu(a_{i,i} + te^{i\theta}) \geq 0, \text{ for all } 0 \leq t < \hat{\rho}_i(\theta),$$

so that the entire complex interval  $[a_{i,i} + te^{i\theta}]_{t=0}^{\hat{\rho}_i(\theta)}$  lies in  $\Gamma^{\mathcal{R}}(A)$ . This implies that the set

$$(2.14) \quad \bigcup_{\theta=0}^{2\pi} [a_{i,i} + te^{i\theta}]_{t=0}^{\hat{\rho}_i(\theta)}$$

is a *star-shaped* subset of  $\Gamma^{\mathcal{R}}(A)$ , for each  $i \in N$ , with

$$(2.15) \quad \nu(a_{i,i} + \hat{\rho}_i(\theta)e^{i\theta}) \in \partial\Gamma^{\mathcal{R}}(A).$$

The results of (2.14) and (2.15) will be used below.

Next, we recall the famous result of Olga Taussky [3], on a sharpening of the Geršgorin Circle Theorem: Let  $A = [a_{i,j}]$  in  $\mathbb{C}^{n,n}$  be irreducible. If  $\lambda \in \sigma(A)$  is such that  $\lambda \notin \text{int } \Gamma_i(A)$  for each  $i \in N$ , i.e.,  $|\lambda - a_{i,i}| \geq r_i(A)$  for each  $i \in N$ , then

$$(2.16) \quad |\lambda - a_{i,i}| = r_i(A), \text{ for each } i \in N,$$

i.e., each Geršgorin circle  $\{z \in \mathbb{C} : |z - a_{i,i}| = r_i(A)\}$  passes through  $\lambda$ .

To complete this section, we include the following:

$$(2.17) \quad \text{If } \nu(z) = 0, \text{ then } \det Q(z) = 0.$$

This follows directly from (2.9), since  $\nu(z)$  is an eigenvalue of  $Q(z)$ . Finally, from [6, Exercise 7, p. 108], we also have that

$$(2.18) \quad \text{for any } z \text{ and } z' \text{ in } \mathbb{C}, |\nu(z) - \nu(z')| \leq |z - z'|,$$

so that  $\nu(z)$  is *uniformly continuous* in  $\mathbb{C}$ . This also will be used below.

**3. Numerical procedure for approximating  $\Gamma^{\mathcal{R}}(A)$ .** With the given irreducible matrix  $A = [a_{ij}]$  in  $\mathbb{C}^{n,n}$ , choose any  $j$  in  $N$ , and set  $z = a_{j,j}$ . Next, we assume that the nonnegative irreducible matrix  $B(a_{j,j})$ , which has at least one zero diagonal entry from (2.3), is a *primitive matrix*; cf. of [5, Section 2.2]. (We note that this is certainly the case if some diagonal entry of  $B(a_{j,j})$  is positive. More generally, if  $B(a_{j,j})$  is not primitive (i.e.,  $B(a_{j,j})$  is cyclic of some index  $p \geq 2$ ), then any simple shift of  $B(a_{j,j})$  into  $B(a_{j,j}) + \varepsilon I_n$  is primitive for each  $\varepsilon > 0$ .)

With  $B(a_{j,j})$  primitive, then, starting with an  $\mathbf{x}^{(0)} > \mathbf{0}$  in  $\mathbb{R}^n$ , the power method gives convergent upper and lower estimates for  $\rho(B(a_{j,j}))$ , i.e., if  $\mathbf{x}^{(m)} := B^m(a_{j,j})\mathbf{x}^{(0)}$  for all  $m \geq 1$ , then with  $\mathbf{x}^{(m)} := [x_1^{(m)}, x_2^{(m)}, \dots, x_n^{(m)}]^T$ , we have

$$(3.1) \quad \underline{\lambda}_m := \min_{i \in N} \left\{ \frac{x_i^{(m+1)}}{x_i^{(m)}} \right\} \leq \rho(B(a_{j,j})) \leq \max_{i \in N} \left\{ \frac{x_i^{(m+1)}}{x_i^{(m)}} \right\} =: \overline{\lambda}_m$$

for all  $m \geq 1$ , with

$$(3.2) \quad \lim_{m \rightarrow \infty} \underline{\lambda}_m = \rho(B(a_{j,j})) = \lim_{m \rightarrow \infty} \overline{\lambda}_m.$$

In this way, using (2.4), (2.7), and (2.9), convergent upper and lower estimates of  $\nu(a_{j,j})$  can be numerically obtained. (These estimations of  $\nu(a_{j,j})$  do not need great accuracy for graphing purposes, as the example in Section 4 shows).

Next, assume, for convenience, that  $\nu(a_{j,j}) > 0$  is accurately known, and select any real  $\theta$ , with  $0 \leq \theta < 2\pi$ . The numerical goal now is to estimate the largest  $\hat{\rho}_j(\theta)$ , with sufficient accuracy, where, from (2.2),

$$(3.3) \quad \nu(a_{j,j} + \hat{\rho}_j(\theta)e^{i\theta}) = 0, \text{ with } \nu(a_{j,j} + (\hat{\rho}_j(\theta) + \varepsilon)e^{i\theta}) < 0$$

for all sufficiently small  $\varepsilon > 0$ . By definition, we then have that

$$(3.4) \quad a_{j,j} + \hat{\rho}_j(\theta)e^{i\theta} \text{ is a boundary point of } \Gamma^{\mathcal{R}}(A).$$

This means, from the min-max conditions (2.8)-(2.9), that there is an  $\mathbf{x} > \mathbf{0}$ , in  $\mathbb{R}^n$ , such that (cf. (2.9))

$$(3.5) \quad Q(a_{j,j} + \hat{\rho}_j(\theta)e^{i\theta})\mathbf{x} = \mathbf{0}, \text{ where } \mathbf{x} = [x_1, x_2, \dots, x_n]^T > \mathbf{0}.$$

Equivalently, on calling  $a_{j,j} + \hat{\rho}_j(\theta)e^{i\theta} =: z_j(\theta)$ , we can express (3.5), using the definition of (2.1), as

$$(3.6) \quad |z_j(\theta) - a_{i,i}| = \sum_{k \in N \setminus \{i\}} |a_{i,k}| x_k / x_i, \quad (\text{all } i \in N),$$

which can be interpreted, from (2.16), simply as Olga Taussky's boundary result. What is perhaps more interesting is that it is geometrically *unnecessary* now to determine the components of the vector  $\mathbf{x} > \mathbf{0}$  in  $\mathbb{R}^n$ , for which (3.6) is valid. This follows since knowing the boundary point  $z_j(\theta)$  of  $\Gamma^{\mathcal{R}}(A)$ , and knowing each of the centers,  $\{a_{i,i}\}_{i \in N}$ , of the associated  $n$  Geršgorin disks, then all the circles of (3.6) can be directly drawn, without knowing the components of the vector  $\mathbf{x}$ .

We return to the numerical estimation of  $\hat{\rho}_j(\theta)$ , which satisfies (3.3)-(3.5). Setting  $z := a_{j,j}$  and  $z' := a_{j,j} + \hat{\rho}_j(\theta)e^{i\theta}$ , we know from (2.18) that

$$(3.7) \quad \hat{\rho}_j(\theta) \geq \nu(a_{j,j}) > 0.$$

Consider then the number  $\nu(a_{j,j} + \nu(a_{j,j})e^{i\theta})$ . If this number is positive, then increase the number  $\nu(a_{j,j})$  to  $\nu(a_{j,j}) + \Delta$ ,  $\Delta > 0$ , until  $\nu(a_{j,j} + (\nu(a_{j,j}) + \Delta)e^{i\theta})$  is negative, and apply a bisection search to the interval  $[\nu(a_{j,j}), \nu(a_{j,j}) + \Delta]$  to determine  $\hat{\rho}_j(\theta)$ , satisfying (3.3). (Again, as in the estimation of  $\nu(a_{j,j})$ , estimates of  $\hat{\rho}_j(\theta)$  do not need great accuracy for graphing purposes.) We remark that a similar bisection search, on  $z$ , can be directly applied to

$$(3.8) \quad \det Q(\nu(a_{j,j} + \hat{\rho}_j(\theta)e^{i\theta})) = 0,$$

as a consequence of (2.11) and (2.15), but this requires, however, the evaluation of an  $n \times n$  determinant.

To summarize, given an irreducible matrix  $A = [a_{i,j}]$  in  $\mathbb{C}^{n,n}$ , our procedure for approximating its minimal Geršgorin set,  $\Gamma^{\mathcal{R}}(A)$ , is to first determine, with reasonable accuracy, the positive numbers  $\{\nu(a_{j,j})\}_{j \in N}$ , and then, again with reasonable accuracy, to determine a few boundary points  $\{\omega_k\}_{k=1}^m$  of  $\Gamma^{\mathcal{R}}(A)$ . For each such boundary point  $\omega_k$  of  $\Gamma^{\mathcal{R}}(A)$ , there is an associated Geršgorin set, consisting of the union of the  $n$  Geršgorin disks, namely,

$$(3.9) \quad \Gamma^{\omega_k}(A) := \bigcup_{i \in N} \{z \in \mathbb{C} : |z - a_{i,i}| \leq |\omega_k - a_{i,i}|\},$$

and their intersection,

$$(3.10) \quad \bigcap_{k=1}^m \Gamma^{\omega_k}(A),$$

gives an approximation to  $\Gamma^{\mathcal{R}}(A)$ , for which  $\Gamma^{\mathcal{R}}(A)$  is a *subset*, and for which  $m$  points, of the boundary of  $\bigcap_{k=1}^m \Gamma^{\omega_k}(A)$ , are *boundary points* of  $\Gamma^{\mathcal{R}}(A)$ .

**4. An easy example.** Consider the irreducible  $3 \times 3$  matrix

$$(4.1) \quad C = \begin{bmatrix} 2 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix},$$

whose minimal Geršgorin set,  $\Gamma^{\mathcal{R}}(C)$ , is shown with the inner blue boundary in Figure 4.1. (This minimal Geršgorin set,  $\Gamma^{\mathcal{R}}(C)$ , also appears as the set with boundary (1) (2) (3) of [6, Figure 4.4].) For the vector  $\mathbf{x}_0 = [1, 1, 1]^T \in \mathbb{R}^3$ , the associated Geršgorin set  $\Gamma^{r^{\mathbf{x}_0}}(C)$ , turns out to be simply

$$(4.2) \quad \Gamma^{r^{\mathbf{x}_0}}(C) = \{z \in \mathbb{C} : |z - 2| \leq 2\}.$$

The boundary of this set is the (outer) *black circle* in Figure 4.1.

Next, starting with the diagonal entry,  $z = 2$ , of the matrix  $C$ , we estimate  $\nu(2)$ , which is positive from (2.12). As  $\mu(2) = 1$  from (2.2), the associated nonnegative irreducible matrix  $B(2)$  from (2.3) is

$$B(2) = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix},$$




FIG. 4.1.

and a few power method iterations (see (3.1)-(3.2)), starting with  $\mathbf{x}_0 = [2, 1, 2]^T$ , gives that  $\rho(B(2)) \doteq 2.2$ . More precisely<sup>1</sup>,  $\rho(B(2)) = 2.24697$ , so that from (2.7) we have  $\nu(2) = 1.24697$ .

Next, we search on the ray  $2 + t$ , with  $t \geq 0$ , for the largest value  $\hat{t}$  for which  $\nu(2 + \hat{t}) = 0$  and  $\nu(2 + t) \geq 0$  for all  $0 \leq t \leq \hat{t}$ . Using the inequality of (2.18), it follows that  $\hat{t} \geq \nu(2) = 1.24697$ . However, in this particular case, it happens that  $\hat{t} = 1.24697$ , so that  $z_1 = 3.24697$  is such that  $\nu(z_1) = 0$ , with  $z_1 \in \partial\Gamma^{\mathcal{R}}(C)$ . Similarly, on considering the diagonal entry  $1 = c_{2,2}$ , we approximate  $\nu(1)$ , which turns out to be  $\nu(1) = 0.80194$ , and then searching on the ray  $1 - t$ ,  $t \geq 0$ , we similarly obtain  $\nu(z_2) = 0$  with  $z_2 = 0.19806$ , and with  $z_2 \in \partial\Gamma^{\mathcal{R}}(C)$ . Calling  $\Gamma^{r_1^{\mathcal{R}}}(C)$  and  $\Gamma^{r_2^{\mathcal{R}}}(C)$  the associated Geršgorin sets, then the intersection of the three sets,  $\bigcap_{j=0}^2 \Gamma^{r_j^{\mathcal{R}}}(C)$ , is shown in Figure 4.1 with the red boundary, where the boundary of the minimal Geršgorin set,  $\Gamma^{\mathcal{R}}(C)$ , is shown in blue.

We see from Figure 4.1 that the set with the red boundary is a set in the complex plane which contains  $\Gamma^{\mathcal{R}}(C)$  and has two real boundary points, shown as the black squares  $z_1$  and  $z_2$ , in common with  $\Gamma^{\mathcal{R}}(C)$ . Continuing, knowing  $\nu(a_{1,1} = a_{3,3} = 2) = 1.24697$  and  $\nu(a_{2,2} = 1) = 0.80194$ , we then look for four additional points of  $\partial\Gamma^{\mathcal{R}}(C)$  which are found on the four rays:  $2 \pm it$ ,  $t \geq 0$ , and  $1 \pm it$ ,  $t \geq 0$ . This gives us the following four points  $\{z_j\}_{j=3}^6$  of  $\Gamma^{\mathcal{R}}(C)$ :

$$z_3 = 1 + i(1.150963), \quad z_4 = \overline{z_3}, \quad z_5 = 2 + i(1.34236), \quad z_6 = \overline{z_5}.$$

The intersection now of the above associated six Geršgorin sets is shown in Figure 4.1 with the green boundary, which includes  $\Gamma^{\mathcal{R}}(C)$  and has six boundary points in common with  $\partial\Gamma^{\mathcal{R}}(C)$ , shown as solid black squares. The region between the green boundary of  $\Gamma^{\mathcal{R}}(C)$  and its blue boundary is colored in yellow, which can be seen as small “roofs” composed of segments of circular arcs.

The amount of numerical calculation to obtain a good approximation to  $\Gamma^{\mathcal{R}}(C)$  is moderate. It is further evident that better approximations to  $\Gamma^{\mathcal{R}}(C)$ , having more points in common with  $\partial\Gamma^{\mathcal{R}}(C)$ , can be similarly constructed.

**5. Comparisons with Braualdi sets.** Given an irreducible matrix  $A = [a_{ij}]$  in  $\mathbb{C}^{n,n}$ ,  $n \geq 2$ , one can similarly associate with  $A$  a minimal Brauer set,  $\mathcal{K}^{\mathcal{R}}(A)$ , as well as a minimal

<sup>1</sup>All such numbers are truncated after five decimal digits.

Brualdi set  $\mathcal{B}^{\mathcal{R}}(A)$ , as described in [6, Section 4.3]. However, it is known (see [6, Theorem 4.15]) that all of these sets are equal, i.e.,

$$(5.1) \quad \Gamma^{\mathcal{R}}(A) = \mathcal{K}^{\mathcal{R}}(A) = \mathcal{B}^{\mathcal{R}}(A),$$

but the approximation of, say, the minimal Brualdi set  $\mathcal{B}^{\mathcal{R}}(A)$ , would now differ from our approximations of the minimal Geršgorin set,  $\Gamma^{\mathcal{R}}(A)$ , described earlier in this paper. For matrices having a very *large* number of nonzero off-diagonal entries, it is *unlikely* (see [6, Section 2.3]) that a similar numerical approximation of the minimal Brualdi set,  $\mathcal{B}^{\mathcal{R}}(A)$ , which from (5.1) equals  $\Gamma^{\mathcal{R}}(A)$ , would be numerically *competitive* with our numerical approach of Section 3 for approximating  $\Gamma^{\mathcal{R}}(A)$ . But, in the case of the matrix  $C$  of (4.1), there are just two associated Brualdi cycles,  $\gamma_1 = (13)$  and  $\gamma_2 = (23)$ , for this matrix  $C$ , so that the approximation of  $\Gamma^{\mathcal{R}}(C)$ , via Brualdi sets, in this case, is easy. In particular, for any  $\mathbf{x} = [x_1, x_2, x_3]^T > \mathbf{0}$  in  $\mathbb{R}^3$ , its associated *Brualdi lemniscates* (cf. [6, eq. (4.78)]) are

$$(5.2) \quad \mathcal{B}_{\gamma_1}^{r^{\mathbf{x}}}(C) = \{z \in \mathbb{C} : |z - 2|^2 \leq r_1^{\mathbf{x}}(C) \cdot r_2^{\mathbf{x}}(C) = \left(\frac{x_3}{x_1}\right) \cdot \left(\frac{x_1 + x_2}{x_3}\right) = \frac{x_1 + x_2}{x_1}\},$$

and

$$(5.3) \quad \mathcal{B}_{\gamma_2}^{r^{\mathbf{x}}}(C) = \{z \in \mathbb{C} : |z - 1||z - 2| \leq \left(\frac{x_3}{x_2}\right) \cdot \left(\frac{x_1 + x_2}{x_3}\right) = \frac{x_1 + x_2}{x_2}\},$$

so that its associated Brualdi set is (cf. [6, eq. (2.40)])

$$(5.4) \quad \mathcal{B}^{r^{\mathbf{x}}}(C) = \mathcal{B}_{\gamma_1}^{r^{\mathbf{x}}}(C) \cup \mathcal{B}_{\gamma_2}^{r^{\mathbf{x}}}(C).$$

Now, knowing that  $z_1 = 3.24697$  is a boundary point of  $\Gamma^{\mathcal{R}}(C)$ , we determine  $\mathbf{x}_1 > \mathbf{0}$  and  $\mathbf{x}_2 > \mathbf{0}$  so that  $z_1 = 3.24697$  is a boundary point of  $\mathcal{B}^{r^{\mathbf{x}_1}}(C)$ . For this particular point  $z_1 = 3.24697$ , the associated Brualdi set, consisting of the union of two Brualdi lemniscate sets, is such that the boundary of *each* Brualdi lemniscate passes through  $z_1$ . (This is exactly the analog of Olga Taussky Theorem in the Geršgorin case; see [1] and [6, Theorem 2.8].) The union of these two Brualdi lemniscate sets can be verified to reduce to

$$\mathcal{B}^{r^{\mathbf{x}_1}}(C) = \{z \in \mathbb{C} : |z - 1| \cdot |z - 2| \leq 2.80193\}.$$

Similarly, for the point  $z_2 = 0.19806$ , the associated Brualdi set has its two lemniscate sets passing through  $z_2$ , and the union of these two Brualdi lemniscate sets can be verified to reduce to the disk

$$\mathcal{B}^{r^{\mathbf{x}_2}}(C) = \{z \in \mathbb{C} : |z - 2| \leq 1.80193\}.$$

The boundary of the intersection  $\mathcal{B}_{\gamma_1}^{r^{\mathbf{x}_1}}(C) \cap \mathcal{B}_{\gamma_2}^{r^{\mathbf{x}_2}}(C)$  is shown in Figure 5.1 with the *green* boundary. Also shown in Figure 5.1, with the *red* boundary, is the related Geršgorin set from Figure 4.1, which also has  $z_1$  and  $z_2$  as common points with the minimal Geršgorin set  $\Gamma^{\mathcal{R}}(C)$ .

From Figure 5.1, we see that  $\mathcal{B}_{\gamma_1}^{r^{\mathbf{x}_1}}(C) \cap \mathcal{B}_{\gamma_2}^{r^{\mathbf{x}_2}}(C)$  is a *proper subset* of the related Geršgorin set, where the difference between these sets is shown in *yellow*. This is not unexpected, as it is known (cf. [6, eq. (4.80)]) that, for any matrix  $A$  in  $\mathbb{C}^{n,n}$ ,

$$\mathcal{B}^{r^{\mathbf{x}}}(A) \subseteq \Gamma^{r^{\mathbf{x}}}(A), \quad \text{for any } \mathbf{x} > \mathbf{0} \text{ in } \mathbb{R}^n.$$




FIG. 5.1.

**Acknowledgment.** This work is supported by the Provincial Secretariat of Science and Technological Development of Vojvodina, Serbia, grant 0708 and by the Ministry of Science of Serbia, grant 144025.

#### REFERENCES

- [1] R. BRUALDI, *Matrices, eigenvalues and directed graphs*, Linear Multilinear Algebra, 11 (1982), pp. 148–165.
- [2] S. GERŠGORIN, *Über die Abgrenzung der Eigenwerte einer Matrix*, Izv. Akad. Nauk SSSR Ser. Mat., 1 (1931), pp. 749–754.
- [3] O. TAUSKY, *Bounds for the characteristic roots of matrices*, Duke Math. J., 15 (1948), pp. 1043–1044.
- [4] R. S. VARGA, *Minimal Geršgorin sets*, Pacific J. Math., 15 (1965), pp. 719–729.
- [5] R. S. VARGA, *Matrix Iterative Analysis*, Second revised and expanded edition, Springer, Berlin, 2000.
- [6] R. S. VARGA, *Geršgorin and His Circles*, Springer, Berlin, 2004.